



Liquid Control Release

Tri-Polymer Fertilizer™

CR K-SUL

AG~NUTRI

Web: www.sjbagnutri.com.au
Email: info@sjbagnutri.com.au
Tech Support: 0407 040 820

ANALYSIS	w/v%
Potassium (as lignosulphonate)	21.0
Sulphur (as sulphate)	11.0

Improve the nitrogen efficiency of tech grade soluble fertilizers by alternating with Control Release Polymers up the dripper.

COMPATIBILITY:

All CR formulations are compatible with each other & through any fertigation system. Please contact SJB Tech Support before mixing CR products with non-CR fertilizers.

RECOMMENDATIONS:

For best results alternate CR products with conventional granular fertilizers such as calcium nitrate. This will improve the nutrient efficacy (so you can lower application rates) of conventional soluble fertilizers as the CR polymer residues in the soil will reduce leaching of nitrogen and other soil mobile nutrients.

CONDITIONS OF SALE:

The Manufacturer warrants that this chelated nutrient product contains the active constituents and polymer/chelating compounds, as specified, and it is for the purpose of supplying the appropriate major nutrient or micro-nutrient, when used in accordance with directions under normal conditions of use. No one, other than the officer of the Manufacturer, is authorised to make any warranty, guarantee or direction concerning the product as the time, place, rate of application and other conditions are beyond the control of the Manufacturer.

Australian Owned & Manufactured by SJB AG~NUTRI PTY. LTD.

©2014

APPLICATION RATES:

General Rate: 10 L/ha every 7-14 days.

Citrus: 10 L/ha at golf ball stage, prior to Autumn flush & at colour change.

Vines: 10 L/ha at fruit set & repeat at bunch closure & again prior to veraison.

For best results alternate CR application with potassium nitrate. After flowering or when crop has optimum leaf area & vigour, cut out soluble potassium nitrate and use CR K-SUL to prevent excessive vigour & promote fruit quality.

GENERAL INFORMATION:

Control Release products contain nutrients in polymer and/or chelate form to provide maximum root absorption when applied through fertigation systems.

SAFETY DIRECTIONS:

Avoid contact with skin and eyes and avoid breathing in vapour or spray mist.

Batch No.	Contents	20L	200L	1000L

