



AG~NUTRI
Pty. Ltd.

Liquid Control Release

Tri-Polymer Fertilizer™



CR NPKomplete

| ANALYSIS | w/v% | | w/v% | | w/v% |
|--------------------------------------|------|--------------------------------|------|--|-------|
| Nitrogen (as polymer) | 10.0 | Magnesium (as lignosulphonate) | 1.0 | Copper (as lignosulphonate) | 0.025 |
| Phosphorous (as ligno-polyphosphate) | 4.0 | Zinc (as lignosulphonate) | 0.1 | Boron (as ammonium ligno-borate) | 0.01 |
| Potassium (as lignosulphonate) | 8.0 | Manganese (as lignosulphonate) | 0.05 | Molybdenum (as ammonium ligno-molybdate) | 0.005 |
| Sulphur (as sulphate) | 3.0 | Iron (as lignosulphonate) | 0.05 | | |

Improve the nutrient efficiency of tech grade soluble fertilizers by alternating with Control Release Polymers up the dripper.

COMPATIBILITY:

Please contact SJB Tech Support before mixing CR products with other CR or conventional fertilizers.

RECOMMENDATIONS:

For best results alternate CR products with conventional granular fertilizers such as calcium nitrate. This will improve the nutrient efficacy (so you can lower application rates) of conventional soluble fertilizers as the CR polymer residues in the soil will reduce leaching of nitrogen and other soil mobile nutrients.

CONDITIONS OF SALE:

The Manufacturer warrants that this chelated nutrient product contains the active constituents and polymer/chelating compounds, as specified, and it is for the purpose of supplying the appropriate major nutrient or micro-nutrient, when used in accordance with directions under normal conditions of use. No one, other than the officer of the Manufacturer, is authorised to make any warranty, guarantee or direction concerning the product as the time, place, rate of application and other conditions are beyond the control of the Manufacturer.

APPLICATION RATES:

General Rate: Early season & post harvest apply 20 L/ha. Maintenance: 10 L/ha every 7-14 days.
Avocado/Mango: 10 L/ha every 14 days during the season. **Capsicum & Tomato:** 20 L/ha at transplanting & 10 L/ha every 7 days after that. **Citrus:** 20 L/ha prior to spring flush & 10 L/ha every month after that. **Cucurbits:** 10 L/ha every 14 days from planting to fruit set. **Strawberries:** 10 L/ha every 7-10 days. **Vines:** 20 L/ha at bud burst & post harvest. During the season apply 10 L/ha every month.

SAFETY DIRECTIONS:

Avoid contact with skin and eyes and avoid breathing in vapour or spray mist.

CONTACT:

Tech Support: 0407 040 820 | info@sjbagnutri.com.au | www.sjbagnutri.com.au

| Batch No. | Contents | 20L | 200L | 1000L |
|-----------|----------|-----|------|-------|
| | | | | |



Australian Owned & Manufactured by SJB AG~NUTRI PTY. LTD. ©2024

