



AG~NUTRI

© SJB AG-NUTRI 2017

LIG-TRACE[®] TAS

For the correction of trace element deficiencies in tree crops, vines, potatoes, poppies & vegetable crops.

**B 1.4%, Cu 0.5%, Mg 1.0%, Mn 2.5%,
Mo 0.4%, S 2.0%, Zn 2.0%**

**Individual Nutrient Chelation Technology
for Maximum Plant Availability.**

www.sjbagnutri.com.au



Australian Owned

APPLICATION

Brassica - Apply 3 L/ha every 14-21 days from early emergence to head formation.

Carrots - Apply 3 L/ha every 4 weeks.

Onions - Apply 2 L/ha at 3 true leaves & 3 L/ha at 4-5 true leaves or prior to bulbing.

Pome Fruit - 2 sprays @ 2-4 L/ha after flowering & post harvest (bud builder).

Poppies - 3 sprays @ 2 L/ha starting prior to hook stage.

Potatoes - 3 L/ha at emergence & repeat prior to row closure.

Stonefruit - 2 L/ha every 10-14 days & 2-4 L/ha post harvest.

Vegetables - Apply 4 L/ha every 14-21 days from early emergence.

Vines - 2-4 L/ha every 14 days from 10cm shoots to fruit set. Repeat after fruit set as required. For young vines apply 2 L/ha every month.



LIG-TRACE TAS is designed for Tasmanian growing conditions for use across a wide range of crops. The unique formulation protects each nutrient & increases foliar uptake, providing superior plant availability over sulphate-based sprays. The high nutrient efficiency means LIG-TRACE TAS can be applied at lower rates per hectare than equivalent non-chelated formulations.

Excellent chemical compatibility

Every nutrient in protected form

Fast & efficient foliar uptake

Low cost per spray

Superior crop safety



SA, TAS, WA & Sunraysia: 0408 633 661, Qld, NT, Northern WA & Northern NSW: 0439 520 744

Technical Support/VIC & Southern NSW: 0407 040 820, E: info@sjbagnutri.com.au

