



AG~NUTRI

www.sjbagnutri.com.au
info@sjbagnutri.com.au
Tech Support: 0407 040 820

Liquid Fully Chelated Lignin Fertilizer™

LIG-ZINC + Boron®

ANALYSIS	w/v%
Zinc (as lignosulphonate)	6.0
Boron (as ammonium ligno-borate)	2.0
Nitrogen (as ligno-ureate)	5.0

Our modified low molecular weight organic chelating agent provides quick absorption & utilization of applied nutrients.

COMPATIBILITY:

This product is compatible with all SJB 'LIG' chelates. With the exception of citrus crops, do not mix with oil sprays. To ensure chelate efficacy it is important that tank water be buffered below pH 7.0. Contact SJB Tech Support before mixing with other nutrient or pesticide sprays.

GENERAL INFORMATION:

This product is chelated (protected from becoming unavailable) with naturally occurring Lignin compounds to provide maximum foliar & root absorption.

CONDITIONS OF SALE:

The Manufacturer warrants that this chelated nutrient product contains the active constituents and chelating compounds, as specified, and it is for the purpose of supplying the appropriate major nutrient or correcting micro-nutrient deficiency, when used in accordance with directions under normal conditions of use. No one, other than the officer of the Manufacturer, is authorised to make any warranty, guarantee or direction concerning the product as the time, place, rate of application and other conditions are beyond the control of the Manufacturer.

Australian Owned & Manufactured by SJB AG~NUTRI PTY. LTD.

©2014

APPLICATION RATES:

General Rate: Maintenance 2 L/ha. Deficiency 4 L/ha.

Almonds: 2 L/ha 10 days after petal fall & 4 L/ha 14 & 28 days later. Young trees - 2-3 L/ha every month. **Apples:** 2 L/ha at budburst & repeat 14 days later. Apply 4 L/ha post-harvest. **Cereal & pulses:** 1-2 L/ha at 5 leaf stage. **Citrus:** 4 L/ha during spring & autumn flushes. **Olives:** 4 L/ha prior to flowering & post harvest. **Potatoes:** 4 L/ha starting at hooking & tuber set. **Vines:** 2 L/ha at 10 cm shoots & repeat 14-21 days later plus 4 L/ha pre-flowering in deficient areas.

RECOMMENDATIONS:

This product may be used as a foliar spray or through fertigation on citrus, stone fruit, grapevines, cereals, pulses, turf, pasture, vegetables, tree crops.

SAFETY DIRECTIONS:

Avoid contact with skin and eyes and avoid breathing in vapour or spray mist.

Batch No.	Contents	20L	200L	1000L

