

Liquid Fully Chelated Lignin Fertilizer Market Mar

LIGNO-BORON®

ANALYSIS	w/v%
Boron (as ammonium ligno-bo	rate) 8.0

Our modified low molecular weight organic chelating agent provides quick absorption & utilization of applied nutrients.

©2017

COMPATIBILITY:

This product is compatible with all SJB 'LIG' chelates. With the exception of citrus crops, do not mix with oil sprays. To ensure chelate efficacy it is important that tank water be buffered below pH 7.0. Contact SJB Tech Support before mixing with other nutrient or pesticide sprays.

GENERAL INFORMATION:

This product is chelated (protected from becoming unavailable) with naturally occurring Lignin compounds to provide maximum foliar & root absorption.

CONDITIONS OF SALE:

The Manufacturer warrants that this chelated nutrient product contains the active constituents and chelating compounds, as specified, and it is for the purpose of supplying the appropriate major nutrient or correcting micronutrient deficiency, when used in accordance with directions under normal conditions of use. No one, other than the officer of the Manufacturer, is authorised to make any warranty, guarantee or direction concerning the product as the time, place, rate of application and other conditions are beyond the control of the Manufacturer.

Australian Owned & Manufactured by SJBAG~NUTRIPTY.LTD.

APPLICATION RATES:

Almonds: Apply 2 L/ha as a pre-bloom bud spray & repeat at fruit set. 4 L/ha post-harvest. Avocado/Mango: 5 L/ha 21 days prior to flowering & repeat every 14-21 days until fruit set. Brassica: 5 L/ha prior to head formation & 2.5 L/ha every 14 days after that. Carrots: 2 L/ha at early & mid-growth stage. Lucerne: 2.5 L/ha prior to flowering Olives: 2.5 L/ha prior to flowering. Poppies: Up to 3 sprays @ 1-2 L/ha from row cover to capsule development. Potatoes: 2 L/ha prior to tuber set & repeat 4 weeks later. Strawberries: 2 L/ha every 14 days from planting to flowering. Vines: 2 L/ha pre-flowering & repeat post-harvest.

RECOMMENDATIONS:

This product may be used as a foliar spray or through fertigation on citrus, stone fruit, grapevines, cereals, pulses, turf, pasture, vegetables, tree crops.

SAFETY DIRECTIONS:

Avoid contact with skin and eyes and avoid breathing in vapour or spray mist.

Batch No.	Contents	20L	200L	1000L
	Contents			

