

Liquid Fully Chelated Lignin Fertilizer

PASTURE CAL

ANALYSIS	w/v%		w/v%
Calcium (as lignosulphonate)	8.0	Nitrogen (as amide)	5.0
Zinc (as lignosulphonate)	0.25	Boron (as ammonium ligno-borate)	0.25

Our modified low molecular weight organic chelating agent provides quick absorption & utilization of applied nutrients.

©2014

COMPATIBILITY:

This product is compatible with all SJB 'LIG' chelates. With the exception of citrus crops, do not mix with oil sprays. To ensure chelate efficacy it is important that tank water be buffered below pH 7.0. Contact SJB Tech Support before mixing with other nutrient or pesticide sprays.

GENERAL INFORMATION:

This product is chelated (protected from becoming unavailable) with naturally occurring Lignin compounds to provide maximum foliar & root absorption.

CONDITIONS OF SALE:

The Manufacturer warrants that this chelated nutrient product contains the active constituents and chelating compounds, as specified, and it is for the purpose of supplying the appropriate major nutrient or correcting micronutrient deficiency, when used in accordance with directions under normal conditions of use. No one, other than the officer of the Manufacturer, is authorised to make any warranty, guarantee or direction concerning the product as the time, place, rate of application and other conditions are beyond the control of the Manufacturer.

Australian Owned & Manufactured by SJB AG~NUTRI PTY. LTD.

APPLICATION RATES:

Lucerne: 5-10 L/ha following a cut or graze, for boost growth add 2-4 L/ha LIG-PASTURE or 5-20 L/ha VLP. Pre plant to early pre emergence 10-100 L/ha. **Dairy Pasture:** 5-10 L/ha pre or early post planting and repeat with 2-4 L/ha LIG-PASTURE or 5-20 L/ha VLP following each grazing for boost growth. Pre plant to early pre-emergence 10-100 L/ha. **Cereal, legumes, canola, crops & fallow:** 10-100 L/ha applied from pre-emergent to 4 leaf crop stage where faster absorption of calcium is required. **DO NOT** mix with Agtryne MA or Igran herbicides.

RECOMMENDATIONS:

This product may be used as a foliar spray or through fertigation on citrus, stone fruit, grapevines, cereals, pulses, turf, pasture, vegetables, tree crops.

SAFETY DIRECTIONS:

Avoid contact with skin and eyes and avoid breathing in vapour or spray mist.

Batch No.	Contents	20L	200L	1000L

