

Liquid Calcium Fertilizer with Added Boron for Enhanced Fruit Quality

ANALYSIS	w/v%
Calcium (as chloride)	16.0
Boron (as boric acid)	0.2

SAFETY DIRECTIONS:

Avoid contact with skin and eves and avoid breathing in vapour or spray mist.

CONDITIONS OF SALE:

The Manufacturer warrants that this nutrient product contains the active constituents, as specified, and it is for the purpose of supplying the appropriate major nutrient or correcting micro-nutrient deficiency, when used in accordance with directions under normal conditions of use. No one, other than the officer of the Manufacturer, is authorised to make any warranty, guarantee or direction concerning the product as the time, place, rate of application and other conditions are beyond the control of the Manufacturer.

> W: www.sjbagnutri.com.au E: info@sibagnutri.com.au Tech Support: 0407 040 820

APPLICATION RATES: 5-10 L/ha every 14-21 days.

The minimum recommended water rate for foliar spraying is 350 L/ha.

WARNING - Application of calcium chloride almost always causes non-serious burn of leaf edges but severe leaf and fruit injury is possible under some conditions. To reduce the chance of severe injury, be careful not to exceed recommended rates. Do not apply the spray under wet, humid, or slow drying conditions. Spraying when temperatures are high may also produce injury. Although the safe upper temperature limit is not known, it is suggested that damage may occur at about 27°C. In hot weather, growers should apply the spray during the coolest part of the day.

Note: StopRotCal B contains a surfactant to enhance product efficacy and therefore additional surfactants (eg. Wetters,

spreaders, stickers) are not required when tank mixing.

Batch No. 201 2001 1000L Contents

Australian Owned & Manufactured by SJB AG~NUTRI PTY. LTD.

©2014